

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

0 579 951 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **93109572.3**

(51) Int. Cl.⁵: **A01N 25/26**

(22) Date of filing: **16.06.93**

(30) Priority: **23.07.92 JP 197228/92**

(43) Date of publication of application:
26.01.94 Patentblatt 94/04

(94) Designated Contracting States:
DE ES FR GR IT

(71) Applicant: **RIKAGAKU KENKYUSHO**
2-1 Hirosawa
Wako-shi Saitama-ken(JP)

(72) Inventor: **Homma, Yasuo, c/o Rikagaku**
Kenkyusho
2-1, Hirosawa
Wako-shi, Saitama-ken(JP)
Inventor: **Arimoto, Yutaka, c/o Rikagaku**
Kenkyusho
2-1, Hirosawa
Wako-shi, Saitama-ken(JP)

(74) Representative: **Werner, Hans-Karsten, Dr. et**
al
Patentanwälte
Von Kreisler-Selting-Werner
Postfach 10 22 41
D-50462 Köln (DE)

(54) **Pesticidal Composition.**

(57) A pesticidal composition comprises a coated pesticide comprising 100 parts by weight of an active pesticidal ingredient coated with 0.1 to 10 parts by weight of one or more ingredients selected from the group consisting of aliphatic acid esters of aliphatic polyols, phospholipids, or a mixture thereof, and 0.01 to 10 parts by weight of a sizing agent. A method for applying a pesticide comprises the step of adding 0.01 to 10 parts by weight of the sizing agent to the coated pesticide, and spraying the resulting mixture.

EP 0 579 951 A1

BACKGROUND OF THE INVENTION

Field of the Invention

- 5 The present invention relates to a pesticidal composition.
 More specifically, the present invention relates to a pesticidal composition which exhibits a prolonged
 pesticidal effect when an active pesticidal ingredient thereof is used at a low concentration.

Description of the Related Art

- 10 Heavy metal compounds such as mercurials and arsenicals, organochlorine agents, and organophosphorus agents have been widely used as agricultural pesticides. However, all of these agents are harmful to human and animal bodies, and furthermore, their use in an effective amount leads to possible environmental pollution, such as soil contamination, which is considered as a serious social problem. For
 15 these reasons, new types of pesticides have been developed which have excellent pesticidal activity and are highly safe to the human body as well as to animals and plants. Japanese Patent Publication No.(Sho)-57-48525 discloses fungicides comprising an aliphatic polyol ester of an aliphatic acid compound and sodium hydrogen carbonate as active ingredients. This publication teaches that the fungicides have high
 20 pesticidal activities against various diseases of plants and stored fruits and that they are very safe to humans, animals and plants. However, in order to obtain sufficient pesticidal effect, it is necessary to use the fungicides at a high concentration when they are sprayed in an amount comparable to those of ordinary pesticides. On the other hand, a large amount of application is necessary when the pesticide is used at a relatively low concentration.

- United States Patent No. 5,123,950 discloses coated agricultural chemicals for eliminating such
 25 problems. These comprise a powdery effective component of an agricultural chemical coated at least with one of aliphatic acid esters of aliphatic polyols and/or phospholipids in an amount of 0.1 to 3 parts by weight per 100 parts by weight of the effective component. Although these coated agricultural chemicals are effective even at a low concentration, they are not satisfactory from the viewpoint of sustained pesticidal activity. Specifically, their prevention values decrease at an advanced stage of disease, although sufficient
 30 pesticidal effect can be achieved at the beginning of their application.

SUMMARY OR THE INVENTION

- Accordingly, an object of the present invention is to provide a pesticidal composition which can exhibit
 35 sustained pesticidal activity even at a low concentration.

- The inventors conducted various studies, and as a result, found that a pesticidal composition comprising a small amount of a sizing agent in addition to a coated pesticide comprising an active pesticidal ingredient coated with 0.1 to 10 parts by weight of one or more ingredients selected from the group consisting of aliphatic acid esters of aliphatic polyols, phospholipids, or a mixture thereof based on 100
 40 parts by weight of the active ingredient can exhibit sustained pesticidal activity at a low concentration. The present invention was achieved on the basis of these findings.

- The present invention thus provides a pesticidal composition which comprises a coated pesticide comprising 100 parts by weight of an active pesticidal ingredient coated with 0.1 to 10 parts by weight of one or more ingredients selected from the group consisting of aliphatic acid esters of aliphatic polyols,
 45 phospholipids, or a mixture thereof, and 0.01 to 10 parts by weight of a sizing agent.

BRIEF EXPLANATION OF THE DRAWINGS

- FIG. 1 shows the results of a comparison of prevention values against citrus melanose using the
 50 pesticidal composition of the present invention (○ - ○), a coated agricultural chemical according to U.S. Patent No. 5,123, 950 (□ - □), and an aqueous NaHCO₃ solution (Δ - Δ).

- FIG. 2 shows the results of a comparison of prevention values against petunia gray mold using the pesticidal composition of the present invention (○ - ○), a coated agricultural chemical according to U.S. Patent No. 5,123, 950 (□ - □), and an aqueous NaHCO₃ solution (Δ - Δ).

55

5

10

25

35

50

ingredient, a coating layer cannot be maintained because of the separation of the aliphatic acid ester of aliphatic polyol and/or the phospholipid, so that an emulsion of the pesticidal composition of the present invention cannot be obtained effectively. The coated pesticides contained in the pesticidal composition of the present invention are generally prepared in the form of powder. Active pesticidal ingredients having a particle size of 100 mesh pass are preferably used, since a coating layer cannot be maintained on particles having too large particle sizes. On the other hand, it is preferable to use active pesticidal ingredients which cannot pass a 400 mesh sieve, since the desired concentration cannot be obtained in the preparation of the coated pesticide where the particles are too small. In general, the coated pesticides are prepared to have a particle size of 100 mesh pass, preferably about 200-400 mesh.

The pesticidal composition of the present invention is characterized in that it contains 0.01-10 parts per weight of a sizing agent together with the above-described coated pesticide. The sizing agent in the form of powder may be mixed with the above-described coated pesticide formulated in the form of powder, or alternatively, the sizing agent may constitute an additional outside coating layer on the coating layer comprising the aliphatic acid ester of aliphatic polyol and/or the phospholipid. Examples of the preparation process of the pesticidal composition of the present invention include, for example, a process of mixing a powdery sizing agent with the powdery coated pesticide; a process of adding the coated pesticide to a solution containing a sizing agent and then evaporating the solvent; and a process comprising the steps of suspending the coated pesticide in a solvent, adding a sizing agent to the resulting suspension, and then evaporating the solvent. As the sizing agent, carboxymethylcellulose (CMC), polyvinylalcohol (PVA), alginic acid, sodium alginate, sodium polyacrylate, sodium pyrophosphate, esters of polyacrylic acid, sodium polyphosphate, potassium polyphosphate, sodium metaphosphate, and synthetic polymer materials may be used. Among the foregoing, CMC is preferably used.

Where the sizing agents in the form of powder are used, 0.01 to 10 parts by weight of powdery sizing agent is mixed with the coated pesticide to prepare the pesticidal composition of the present invention. In this case, powdery sizing agent having a particle size of 325 mesh pass can be added and mixed with the coated pesticide having a particle size of 100 mesh pass, preferably about 200 to 400 mesh pass, in the ratio of 0.01 to 10 parts by weight based on 100 parts by weight of active pesticidal ingredient. Alternatively, the pesticidal composition of the present invention in which the sizing agent is coated on the outside of the coated pesticide can be prepared by adding the coated pesticide to the solution prepared by dissolving the sizing agent in water or an organic solvent, and then evaporating the solvent. Where the sizing agent is rather insoluble in water, organic solvents which can dissolve the sizing agent are preferably used. Preferably, these organic solvents do not substantially affect the coating layer of the aliphatic acid ester of aliphatic polyol and/or the phospholipid during the coating with the sizing agent. Such solvents can appropriately be chosen by one of ordinary skill in the art and examples thereof include ethanol and methanol. Furthermore, the pesticidal composition of the present invention can be prepared by dissolving the sizing agent in a suspension prepared by suspending the coated pesticide in water or an organic solvent, and then evaporating the solvent. As the solvent, there may be chosen one that can dissolve the sizing agent and but does not substantially affect the coating layer of the aliphatic acid ester of aliphatic polyol and/or the phospholipid during the coating process with the sizing agent. For example, water, methanol, or ethanol may be used.

The pesticidal composition of the present invention may be formulated by optionally adding commonly used auxiliary agents such as spreading agents, wetting and spreading agents, fixing agents and surfactants. These auxiliary agents in the form of powder may be added to the pesticidal composition of the present invention, or alternatively, they may be contained in the coated pesticide or the coating of the sizing agent. The concentrations for spraying the pesticidal composition of the present invention are not limited since they may vary depending on the sort of active ingredient. For example, they may be used at about 1 to 500 ppm.

The present invention also provides a method for applying a pesticide comprising the step of adding 0.01 to 10 parts by weight of a sizing agent to the coated pesticide prepared by the method described above, and spraying the resulting mixture. For example, where the coated pesticide is used as a wettable powder, the process may comprise the steps of dissolving the above-described sizing agent in the aqueous preparation of the coated pesticide having a concentration of about 100 to 1000 ppm so that the content ratio of the sizing agent is 0.01 to 10 parts by weight based on 100 parts by weight of the active pesticidal ingredient, and spraying the resulting mixture. While the invention is not bound by any theory, it is thought that when a spray application is carried out according to the present invention, a highly concentrated solution of an active pesticide is kept in a capsule formed by the aliphatic acid ester of aliphatic polyol and/or the phospholipid since the sizing agent and the active pesticidal ingredient contained in the coated pesticide are readily dissolved in water while the aliphatic acid ester of aliphatic polyol and/or the

phospholipid are water-insoluble. On the other hand, the outside of the capsule is contacted with dissolved sizing agent, and as a consequence, the capsule containing the highly concentrated pesticidal solution will readily adhere to pathogenic microorganisms or injurious insects and exhibit sustained bactericidal and insecticidal effects. Where the spray application is carried out using the pesticidal compositions of the present invention added with water, they can exhibit sustained bactericidal and insecticidal effect in a similar manner even at a low concentration.

EXAMPLES

The present invention will be hereinafter explained more specifically by way of examples. However, the present invention is not be limited to these examples.

Example 1

Diglycerin laurate (2 g) was dissolved in 100 ml of acetone. To the resulting solution, 100 g of finely powdered sodium hydrogen carbonate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of carboxymethylcellulose (CMC) was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 2

Glycerin monocaprylate (4 g) was dissolved in 100 ml of acetone. To the resulting solution, 100 g of finely powdered sodium hydrogen carbonate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.02 g of carboxymethylcellulose (CMC) was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 3

Glycerin monolaurate (2 g) and sorbitan stearate (2 g) were dissolved in 100 ml of ethanol. To the resulting solution, 100 g of finely powdered sodium chloride (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.1 g of carboxymethylcellulose (CMC) was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 4

Glycerin monocaprylate (2 g) and diglycerin laurate (2 g) were dissolved in 100 ml of ethanol. To the resulting solution, 100 g of finely powdered potassium hydrogen carbonate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium polyacrylate was added and mixed as the sizing agent to give powdery water soluble concentrate having good fluidity.

Example 5

Glycerin monolaurate (4 g) and diglycerin oleate (2 g) were dissolved in 10 ml of acetone and 5 ml of methanol, respectively. To the solution of glycerin monolaurate, 100 g of finely powdered potassium carbonate (90% 100 mesh pass) was added and the mixture was hot-air dried after vigorous stirring. The solution of diglycerin oleate was added to the result and vigorously stirred, and the solvent was then thoroughly removed by hot-air drying. To the resulting powder, 0.08 g of sodium polyacrylate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 6

Glycerin monooleate (2 g) and diglycerin oleate (2 g) were dissolved in 50 ml of acetone and 50 ml of methanol, respectively. To the solution of glycerin monooleate, 100 g of finely powdered ammonium hydrogen carbonate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly

removed using a rotary evaporator. The solution of diglycerin oleate was added to the result and vigorously stirred, and the solvent was then thoroughly removed by hot-air drying. To the resulting powder, 0.05 g of sodium alginate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 7

Glycerin laurate (4 g) was dissolved in 100 ml of methanol: To the resulting solution, 80 g of finely powdered ammonium carbonate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of alginic acid was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 8

Diglycerin oleate (2 g) and hexaglycerin laurate (2 g) were dissolved in 50 ml of acetone and 50 ml of methanol, respectively. To the solution of diglycerin oleate, 100 g of finely powdered potassium carbonate (90% 100 mesh pass) was added and the solvent was thoroughly removed using a rotary evaporator after vigorous stirring. The solution of hexaglycerin laurate was added to the result and vigorously stirred, and the solvent was then thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium pyrophosphate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 9

Diglycerin laurate (2 g) and decaglycerin laurate (2 g) were dissolved in 50 ml of ethanol and 50 ml of acetone, respectively. To the former solution, 100 g of finely powdered disodium phosphate (90% 100 mesh pass) was added and the solvent was thoroughly removed using a rotary evaporator after vigorous stirring. The solution of decaglycerin laurate was added to the result and vigorously stirred, and the solvent was then thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium polyphosphate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 10

Propyleneglycol monolaurate (4 g) and sorbitan laurate (2 g) were dissolved in 100 ml of ethanol. To the resulting solution, 100 g of finely powdered monosodium phosphate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium polyphosphate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 11

Diglycerin laurate (2 g) and decaglycerin laurate (2 g) were dissolved in 50 ml of ethanol and 50 ml of acetone, respectively. To the solution of diglycerin laurate, 100 g of finely powdered dipotassium phosphate (90% 100 mesh pass) was added and the solvent was thoroughly removed using a rotary evaporator after vigorous stirring. The solution of decaglycerin laurate was added to the result and vigorously stirred, and the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium polyphosphate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 12

Propyleneglycol monolaurate (4 g) and sorbitan laurate (2 g) were dissolved in 100 ml of ethanol. To the resulting solution, 100 g of finely powdered monopotassium phosphate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of sodium polyphosphate was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Example 13

Glycerin monooleate (4 g) was dissolved in 100 ml of acetone. To the resulting solution, 100 g of finely powdered anhydrous copper sulfate (90% 100 mesh pass) was added. After vigorous stirring, the solvent was thoroughly removed using a rotary evaporator. To the resulting powder, 0.05 g of alginic acid was added and mixed as the sizing agent to give a powdery water soluble concentrate having good fluidity.

Experiments 1-13

The efficacy of the pesticidal compositions of the present invention was compared with that of the coated agricultural chemicals disclosed in U.S. Patent 5,123,950 and the solutions of commonly used active ingredients. Pesticidal formulations were prepared at the concentrations of 250 ppm and 125 ppm for each pesticide. Comparison was made by calculating prevention values (%) at 3, 6, and 10 days after the application and estimating the presence or absence of phytotoxicity. In Table 1, each coated agricultural chemical comprises the same ingredients as the corresponding composition of the present invention except that each coated agricultural chemical does not contain a sizing agent. The preventive values of the 125 ppm formulations are shown in parentheses in the table, and their phytotoxicities are not shown since the same results were obtained by the formulations at the concentration of 250 ppm. The result obtained by the application of a formulation not containing an active ingredient is shown for comparison.

The results summarized in Table 1 demonstrate that the pesticidal composition of the present invention exhibits markedly higher preventive values even at low concentrations of 125 ppm and 250 ppm as compared with ordinarily used pesticidal solutions or known coated agricultural chemicals and that they achieve excellent preventive values at 6 and 10 days and have sustained pesticidal activity.

Table 1

Efficacy of the Treatment for Cucumber Powdery Mildew

(*Sphaerotheca fuliginea*)

| Pesticide Tested (sizing agent) | Efficacy (%) 3, 6, and 10 days | | | Phyto- toxicity |
|---|-----------------------------------|----------------------------|----------------------------|--------------------|
| Example 1 (CMC) Coated Agricultural Chemical NaHCO ₃ | 95(87) 90(55) 30(20) | 93(85) 85(51) 27(17) | 93(80) 80(48) 20(10) | N.D. " " |
| Example 2 (CMC) Coated Agricultural Chemical Na ₂ CO ₃ | 90(48) 89(48) 14(5) | 86(45) 84(46) 12(0) | 85(45) 45(27) 5(0) | " " " |
| Example 3 (CMC) Coated Agricultural Chemical NaCl | 85(50) 90(45) 10(0) | 80(45) 85(41) 9(0) | 65(27) 41(18) 5(0) | " " " |
| Example 4 (sodium polyacrylate) Coated Agricultural Chemical KHCO ₃ | 96(89) 98(58) 12(10) | 93(87) 92(57) 10(5) | 91(87) 57(50) 5(0) | " " " |
| Example 5 (sodium polyacrylate) Coated Agricultural Chemical K ₂ CO ₃ | 80(45) 91(45) 10(5) | 79(40) 88(42) 10(5) | 79(40) 42(20) 5(0) | " " " |
| Example 6 (alginic acid) Coated Agricultural Chemical (NH ₄)HCO ₃ | 93(70) 83(64) 12(5) | 91(66) 80(53) 9(0) | 90(66) 53(40) 0(0) | " " " |
| Example 7 (alginic acid) Coated Agricultural Chemical (NH ₄) ₂ CO ₃ | 88(66) 86(52) 9(0) | 86(61) 82(51) 7(0) | 85(62) 51(35) 0(0) | " " " |
| Example 8 (sodium pyrophosphate) Coated Agricultural Chemical CaCO ₃ | 86(50) 87(50) 7(0) | 85(48) 81(47) 5(0) | 81(48) 47(27) 0(0) | " " " |
| Example 9 (sodium polyphosphate) Coated Agricultural Chemical Na ₂ HPO ₄ | 90(80) 90(51) 14(6) | 88(71) 81(50) 10(3) | 87(70) 50(25) 5(0) | " " " |
| Example 10 (sodium polyphosphate) Coated Agricultural Chemical NaH ₂ PO ₄ | 86(50) 100(50) 5(0) | 82(44) 93(46) 0(0) | 83(44) 46(18) 0(0) | " " " |

Table 1 (Continued)

| Pesticide Tested (sizing agent) | Efficacy (%) 3, 6, and 10 days | | | Phyto- toxicity |
|--|-----------------------------------|----------------------------|----------------------------|--------------------|
| Example 11 (sodium polyphosphate) Coated Agricultural Chemical K ₂ HPO ₄ | 79(44) 94(51) 6(0) | 75(40) 93(49) 5(0) | 75(40) 49(20) 3(0) | N.D. " " |
| Example 12 (sodium polyphosphate) Coated Agricultural Chemical KH ₂ PO ₄ | 83(55) 90(48) 11(5) | 80(51) 89(46) 7(0) | 80(51) 46(20) 3(0) | " " " |
| Example 13 (alginic acid) Coated Agricultural Chemical CuSO ₄ | 95(77) 90(75) 19(15) | 95(70) 88(70) 17(13) | 92(70) 52(34) 15(13) | " " " |
| Water (Reference) | 0 | 0 | 0 | " |

N.D.: not detected

Experiment 14

The efficacy of the treatment was studied using the compositions of the present invention in the form of water soluble concentrates, each containing a different sizing agent. Pesticidal formulations were prepared at the concentrations of 250 ppm and 125 ppm for each pesticide. Comparison was made by calculating preventive values (%) at 3, 6, and 10 days after the application and estimating the presence or absence of phytotoxicity. In Table 1, the preventive values of the 125 ppm formulations are shown in parentheses and their phytotoxicities are not shown since the same results were obtained by the formulations at the concentration of 250 ppm. The water soluble concentrate of the present invention had successful efficacy and sustained activity irrespective the sort of sizing agent contained therein. The results are summarized in Table 2.

Table 2

Efficacy of the Treatment for Cucumber Powdery Mildew

| Coated Pesticide/sizing agent | Efficacy (%) 3, 6, and 10 days | | | Phyto- toxicity |
|--------------------------------|-----------------------------------|--------|--------|--------------------|
| Coated NaHCO ₃ /CMC | 96(85) | 96(85) | 94(83) | N.D. |
| " /alginic acid | 95(85) | 94(85) | 94(82) | " |
| " /casein | 90(81) | 90(82) | 90(80) | " |
| " /sodium polyacrylate | 95(87) | 95(86) | 95(86) | " |
| " /sodium pyrophosphate | 94(82) | 93(80) | 93(79) | " |
| " /sodium polyphosphate | 93(80) | 91(80) | 91(77) | " |
| " /sodium metaphosphate | 80(70) | 80(69) | 77(65) | " |
| Coated NaHCO ₃ / — | 64(51) | 87(62) | 50(49) | " |

Experiment 15

Fig. 1 shows the results of a comparison in preventive values against citrus melanose in which an aqueous NaHCO₃ solution (Δ - Δ), the dilution of the coated agricultural chemical according to U.S. Patent No. 5,123,950 (\square - \square), and the dilution of the pesticidal composition of the present invention (O - O) were used at various concentrations ranging from 0 to 300 ppm. The preventive value of the ordinarily used pesticidal solution was 5% even at 300 ppm, and the preventive value of the the coated agricultural chemical according to U.S. Patent No. 5,123,950 was about 60% at 250 ppm. On the other hand, the pesticidal composition of the present invention exhibited a preventive value of not lower than 80% at 125 ppm. The pesticidal composition of the present invention is obviously far more effective at low concentration than the other two pesticides.

Experiment 16

FIG. 2 shows the results of a comparison in preventive values against petunia gray mold (*Botrytis cinerea*) in which an aqueous NaHCO₃ solution (Δ - Δ), the dilution of the coated agricultural chemical according to U.S. Patent No. 5,123,950 (\square - \square), and the dilution of the pesticidal composition of the present invention (O - O) were used. It is apparent that the pesticidal composition of the present invention is far more effective at low concentration than the other two pesticides.

Experiment 17

The insecticidal activity of the solution of the pesticidal composition of the present invention was compared with those of known coated agricultural chemical according to U.S. Patent No. 5,123,950 and ordinarily used pesticidal solution. Formulations at 250 ppm and 125 ppm were prepared for each of these pesticides. Insecticidal rate (%) against *Myzus persicae* (green peach aphid) and mitecidal rate (%) against *Panonychus citri* McGregor (citrus red mite) were calculated for comparison. In the inhibitory activity against *Icerya purchasi* Maskell (cottony cushion scale) shown in Table 3, + represents effective; ++ represents highly effective; and - represents ineffective. The results the 125 ppm formulations are shown in parentheses in the table.

From these results, it is apparent that the aqueous solution of the compositions of the present invention have higher insecticidal activities than ordinarily used pesticidal solution and known coated agricultural chemical according to U.S. Patent No. 5,123,950. In particular, as compared with other pesticides, it is apparent that remarkable results were obtained by the pesticidal compositions of Examples 1 to 13 for

insecticidal rate (%) against *Myzus persicae*; by the pesticidal compositions of Examples 1, 4, 6, 8, 10 and 11 for mitecidal rate (%) against *Panonychus citri* MGregor; and by the pesticidal composition of Examples 1, 4, 6, 11, and 13 for inhibitory activity against *Icerya purchasi* Maskell. The results are summarized in Table 3.

Table 3

Insecticidal and Mitecidal Effect

| Pesticide Tested (sizing agent) | Insecticidal rate ^a | Mitecidal rate ^b | Inhibitory effect ^c |
|---|-----------------------------------|--------------------------------|-----------------------------------|
| Example 1 (CMC) | 90(55) | 97(62) | ++(++) |
| Coated Agricultural Chemical | 75(31) | 88(46) | +(-) |
| NaHCO ₃ | 20 (0) | 27(10) | -(-) |
| Example 2 (CMC) | 98(51) | | |
| Coated Agricultural Chemical | 58(23) | | |
| Na ₂ CO ₃ | 10 (0) | | |
| Example 3 (CMC) | 83(49) | | |
| Coated Agricultural Chemical | 55(26) | | |
| NaCl | 17(10) | | |
| Example 4 (Na polyacrylate) | 100(88) | 98(84) | ++(++) |
| Coated Agricultural Chemical | 89(52) | 95(61) | ++(+) |
| KHCO ₃ | 20(15) | 50(20) | -(-) |
| Example 5 (Na polyacrylate) | 96(50) | | |
| Coated Agricultural Chemical | 80(31) | | |
| K ₂ CO ₃ | 27(11) | | |
| Example 6 (alginic acid) | 90(44) | 97(73) | ++(++) |
| Coated Agricultural Chemical | 81(25) | 88(48) | ++(+) |
| (NH ₄)HCO ₃ | 20(11) | 20(7) | -(-) |
| Example 7 (alginic acid) | 94(59) | | |
| Coated Agricultural Chemical | 79(48) | | |
| (NH ₄) ₂ CO ₃ | 10 (0) | | |
| Example 8 (Na pyrophosphate) | 72(40) | 90(71) | |
| Coated Agricultural Chemical | 30(18) | 83(40) | |
| CaCO ₃ | 0 (0) | 21 (7) | |
| Example 9 (Na polyphosphate) | 97(62) | | |
| Coated Agricultural Chemical | 77(35) | | |
| Na ₂ HPO ₄ | 10 (0) | | |
| Example 10 (Na polyphosphate) | 91(72) | 93(87) | |
| Coated Agricultural Chemical | 70(33) | 76(42) | |
| NaH ₂ PO ₄ | 11 (0) | 22 (7) | |

Table 3 (Continued)

| Pesticide Tested (sizing agent) | Insecticidal rate ^a | Mitecidal rate ^b | Inhibitory effect ^c |
|------------------------------------|-----------------------------------|--------------------------------|-----------------------------------|
| Example 11 (Na polyphosphate) | 95(66) | 92(80) | ++(++) |
| Coated Agricultural Chemical | 77(37) | 75(32) | +(-) |
| K ₂ HPO ₄ | 12 (7) | 20(11) | -(-) |
| Example 12 (Na polyphosphate) | 93(71) | | |
| Coated Agricultural Chemical | 68(38) | | |
| KH ₂ PO ₄ | 15(10) | | |
| Example 13 (alginic acid) | 100(65) | | ++(++) |
| Coated Agricultural Chemical | 80(32) | | ++(-) |
| CuSO ₄ | 36(12) | | -(-) |
| Water (Reference) | 0 | 0 | -(-) |

a: Insecticidal rate (%) against Myzus persicae

b: Mitecidal rate (%) against Panonychus citri McGregor

c: Inhibitory activity against Icerya purchst Maskell

Claims

1. A pesticidal composition which comprises a coated pesticide comprising 100 parts by weight of an active pesticidal ingredient coated with 0.1 to 10 parts by weight of one or more ingredients selected from the group consisting of aliphatic acid esters of aliphatic polyols, phospholipids, or the mixture thereof, and 0.01 to 10 parts by weight of a sizing agent.
2. The pesticidal composition according to claim 1, wherein the aliphatic acid ester of aliphatic polyol is selected from the group consisting of sorbitan monolaurate, sorbitan monostearate, glycerin laurate, glycerin monocaprylate, glycerin monooleate, glycerin monooctate, glycerin mono-soyabean oil fatty acid ester, triglycerin monooleate, glycerin monopalmitate, and polyglycerin fatty acid esters, polyoxyethylene-alkylphenylether, polyoxyethylene-nonylphenylether, lauryl dihydroxyethylamine, and mixtures thereof.
3. The pesticidal composition according to claim 1, wherein the phospholipid is selected from the group consisting of purified lecithin, phosphatidylethanolamine, phosphatidylinositol, and mixtures thereof.
4. The pesticidal composition according to claim 1, wherein the sizing agent is selected from the group consisting of carboxymethylcellulose, polyvinylalcohol, alginic acid, sodium alginate, sodium polyacrylate, sodium pyrophosphate, esters of polyacrylic acid, sodium polyphosphate, potassium polyphosphate, sodium metaphosphate, and mixtures thereof.
5. A method for applying a pesticide comprising the step of adding 0.01 to 10 parts by weight of a sizing agent to a coated pesticide comprising 100 parts by weight of an active pesticidal ingredient coated with 0.1 to 10 parts by weight of one or more ingredients selected from the group consisting of aliphatic acid esters of aliphatic polyols, phospholipids, or a mixture thereof, and spraying the resulting mixture.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 10 9572

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|--|---|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.Cl.5) |
| Y | FR-A-2 623 973 (RIKAGAKU KENKYUSHO) * page 6, line 9 - line 15 * * page 6, line 35 - line 38 * D & US-A-5 123 950 --- | 1-5 | A01N25/26 |
| Y | DE-A-29 27 994 (RIKAGAKU KENKYUSHO) * claims 1-4 * * page 14, paragraph 2 -paragraph 3 * D & JP-B-57 048 525 (...) --- | 1-5 | |
| Y | GB-A-2 002 319 (BATELLE MEMORIAL INSTITUTE) * the whole document * ----- | 1-5 | |
| | | | TECHNICAL FIELDS SEARCHED (Int.Cl.5) |
| | | | A01N |
| The present search report has been drawn up for all claims | | | |
| Place of search THE HAGUE | | Date of completion of the search 9 November 1993 | Examiner LAMERS, W |
| CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document | | T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document | |